A&A manuscript no.
(will be inserted by hand later)

Your thesaurus codes are:
11(11.09.1 NGC 6946; 11.09.4, 11.19.2, 11.19.6), 13(13.09.1, 13.09.3)

ASTRONOMY
AND
ASTROPHYSICS
21.10.1996

ISOCAM* Observations of NGC 6946: Mid-IR Structure

S. Malhotra,[1], G. Helou[1], D. Van Buren[1], M. Kong[1], C. A. Beichman[1], H. Dinerstein[2], D. J. Hollenbach[3], D. A. Hunter[4], K. Y. Lo[5], S. D. Lord[1], N. Y. Lu[1], R. H. Rubin[3], G. J. Stacey[6], H. A. Thronson Jr.[7], and M. W. Werner[8]

[1] IPAC, California Institute of Technology, MS 100-22, Pasadena, CA 91125
[2] University of Texas, Astronomy Department, RLM 15.308, Texas, Austin, TX 78712
[3] NASA/Ames Research Center, MS 245-6, Moffett Field, CA 94035
[4] Lowell Observatory, 1400 Mars Hill Rd., Flagstaff, AZ 86001
[5] University of Illinois, Astronomy Department, 1002 W. Green St., Urbana, IL 61801
[6] Cornell University, Astronomy Department, 220 Space Science Building, Ithaca, NY 14853
[7] University of Wyoming, Wyoming Infrared Observatory, Laramie, WY, 82071
[8] Jet Propulsion Laboratory, MS 233-303, 4800 Oak Grove Rd., Pasadena, CA 91109

Received August 7 1996; accepted 19th August

Abstract. The nearby spiral galaxy NGC 6946 was observed with ISO-CAM in the mid-infrared, achieving 7″ resolution and sub- MJy sr^{-1} sensitivity. Images taken with CAM filters LW2 (7μm) and LW3 (15μm) are analysed to determine the morphology of this galaxy and understand better the emission mechanisms. The mid-infrared emission follows an exponential disk with a scale length 75″. This is 60 % of the scale length in the optical R-band and radio continuum. The nuclear starburst region is too bright for reliable measurement in these images. Its surface brightness exceeds the inner disk by at least a factor of 12. The arms and interarm regions are clearly detected, with each of these components contributing about equally to the disk emission. The arm-interarm contrast is 2-4 in the mid-IR, close to that measured in the visible R band light and lower than the contrast in $H\alpha$, suggesting that non-ionizing radiation contributes significantly to dust heating.

Key words: Infrared:galaxies – Infrared:ISM:continuum – Galaxies:ISM – Galaxies:spiral – Galaxies:structure

1. Introduction

NGC 6946 is a nearby face-on spiral galaxy, a good target for a study of star-formation and related processes in "normal" galaxies. In addition to the well-resolved disk, it has a starburst nucleus (*e.g.* Engelbracht et al. 1996), and a bright northern spiral arm that earned it an entry in the Atlas of Peculiar Galaxies (Arp 1966). NGC 6946 was therefore selected as a focus of the Key Project on the interstellar medium of normal galaxies (Helou et al. 1996) under NASA Guaranteed Time on ISO (Kessler et al. 1996). We report here on the ISO-CAM (Césarsky et al. 1996) maps obtained for this galaxy, revealing for the first time the mid-infrared (mid-IR) morphology at vastly improved resolution and sensitivity.

Mid-IR emission in the 5 to 20 μm range is dominated by very small grains fluctuating to high temperatures and from Poly Aromatic Hydrocarbons (PAH) features. Even though these grains are not in thermal equilibrium, they still convert heating photons (Draine and Anderson 1985), and should therefore trace star-formation, allowing a detailed and un-extincted view of that activity. By studying the distribution of the mid-IR emission with respect to other components of the galaxy, i.e. HI, H_2, ionized gas, and starlight we can also characterize the heating sources for the mid-IR, and better define its diagnostic value.

2. Observations and Data reduction

NGC 6946 was mapped at 7μm (LW2 filter, $\Delta\lambda = 3.5\mu$m) and at 15 μm (LW3 filter, $\Delta\lambda = 6\mu$m), using the raster scan mode to cover roughly $12.5' \times 12.5'$ centered on the nucleus. CAM was set to 6″/pixel, and the raster was made up of 8×8 pointings separated by 81″, or 13.5 pixels in each direction, allowing for better spatial sampling. At each pointing in the raster scan we took 8 frames integrating 5 seconds for each at 7 μm, and 10 frames of 2 seconds each at 15 μm. Hα and broad-band imaging

Send offprint requests to: san@ipac.caltech.edu
* Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) and with the participation of ISAS and NASA.